1 Family Planning

Mr. and Mrs. Brown decide to continue having children until they either have their first girl or until they have three children. Assume that each child is equally likely to be a boy or a girl, independent of all other children, and that there are no multiple births. Let *G* denote the numbers of girls that the Browns have. Let *C* be the total number of children they have.

- (a) Determine the sample space, along with the probability of each sample point.
- (b) Compute the joint distribution of *G* and *C*. Fill in the table below.

	C=1	C=2	C=3
G=0			
G=1			

(c) Use the joint distribution to compute the marginal distributions of *G* and *C* and confirm that the values are as you'd expect. Fill in the tables below.

Pr(G=0)	
Pr(G=1)	

Pr(C=1)	Pr(C=2)	Pr(C=3)

- (d) Are *G* and *C* independent?
- (e) What is the expected number of girls the Browns will have? What is the expected number of children that the Browns will have?

2 Linearity

Solve each of the following problems using linearity of expectation. Explain your methods clearly.

- (a) In an arcade, you play game A 10 times and game B 20 times. Each time you play game A, you win with probability 1/3 (independently of the other times), and if you win you get 3 tickets (redeemable for prizes), and if you lose you get 0 tickets. Game B is similar, but you win with probability 1/5, and if you win you get 4 tickets. What is the expected total number of tickets you receive?
- (b) A monkey types at a 26-letter keyboard with one key corresponding to each of the lower-case English letters. Each keystroke is chosen independently and uniformly at random from the 26 possibilities. If the monkey types 1 million letters, what is the expected number of times the sequence "book" appears?
- (c) A building has n floors numbered 1, 2, ..., n, plus a ground floor G. At the ground floor, m people get on the elevator together, and each gets off at a uniformly random one of the n floors (independently of everybody else). What is the expected number of floors the elevator stops at (not counting the ground floor)?
- (d) A coin with heads probability p is flipped n times. A "run" is a maximal sequence of consecutive flips that are all the same. (Thus, for example, the sequence HTHHHTTH with n=8 has five runs.) Show that the expected number of runs is 1+2(n-1)p(1-p). Justify your calculation carefully.

3 Coupon Collection

Suppose you take a deck of n cards and repeatedly perform the following step: take the current top card and put it back in the deck at a uniformly random position. (The probability that the card is placed in any of the n possible positions in the deck — including back on top — is 1/n.) Consider the card that starts off on the bottom of the deck. What is the expected number of steps until this card rises to the top of the deck? (*Hint*: Let T be the number of steps until the card rises to the top. We have $T = T_n + T_{n-1} + \cdots + T_2$, where the random variable T_i is the number of steps until the bottom card rises from position i to position i-1. Thus, for example, T_n is the number of steps until the bottom card rises off the bottom of the deck, and T_2 is the number of steps until the bottom card rises from second position to top position. What is the distribution of T_i ?)